


# o8: Poisson and More

---



Jerry Cain  
April 17<sup>th</sup>, 2024

[Lecture Discussion on Ed](#)

# Poisson



# Algorithmic ride sharing



Probability of  $k$  requests from this area in the next 1 minute?

Suppose we know:

On average,  $\lambda = 5$  requests per minute

# Algorithmic ride sharing, approximately

Probability of  $k$  requests from this area in the next 1 minute?

On average,  $\lambda = 5$  requests per minute

Break a minute down into 60 seconds:



At each second:

- Independent Bernoulli trial
- You get a request (1) or you don't (0).

Let  $X = \#$  of requests in minute.

$$E[X] = \lambda = 5$$

$$X \sim \text{Bin}(n = 60, p = 5/60)$$

$$P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$$




But what if there are *two* requests in the same second?

# Algorithmic ride sharing, approximately

Probability of  $k$  requests from this area in the next 1 minute?

On average,  $\lambda = 5$  requests per minute

Break a minute down into 60,000 milliseconds:



At each millisecond:

- Independent Bernoulli trial
- You get a request (1) or you don't (0).

Let  $X = \#$  of requests in minute.

$$E[X] = \lambda = 5$$

$$X \sim \text{Bin}(n = 60000, p = \lambda/n)$$

$$P(X = k) = \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k}$$




But what if there are *two* requests in the same millisecond?

# Algorithmic ride sharing, approximately

Probability of  $k$  requests from this area in the next 1 minute?

On average,  $\lambda = 5$  requests per minute

Break a minute down into **infinitely small** buckets:



For each time bucket:

- Independent Bernoulli trial
- You get a request (1) or you don't (0).

Let  $X = \#$  of requests in minute.

$$E[X] = \lambda = 5$$

$$X \sim \text{Bin}(n, p = \lambda/n)$$

$$P(X = k) = \lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k}$$

Gnarly math incoming!

# Binomial in the limit

$$\lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^n = e^{-\lambda}$$


$$P(X = k) = \lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} \stackrel{\text{Expand}}{=} \lim_{n \rightarrow \infty} \frac{n!}{k!(n-k)!} \frac{\lambda^k}{n^k} \frac{\left(1 - \frac{\lambda}{n}\right)^n}{\left(1 - \frac{\lambda}{n}\right)^k}$$

$$\stackrel{\text{Rearrange}}{=} \lim_{n \rightarrow \infty} \frac{n!}{n^k(n-k)!} \frac{\lambda^k}{k!} \frac{\left(1 - \frac{\lambda}{n}\right)^n}{\left(1 - \frac{\lambda}{n}\right)^k} \stackrel{\text{Def natural exponent}}{=} \lim_{n \rightarrow \infty} \frac{n!}{n^k(n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

$$\stackrel{\text{Expand}}{=} \lim_{n \rightarrow \infty} \frac{n(n-1)\cdots(n-k+1)}{n^k} \frac{(n-k)!}{(n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

$$\stackrel{\text{Limit analysis + cancel}}{=} \lim_{n \rightarrow \infty} \frac{n^k}{n^k} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{1} \stackrel{\text{Simplify}}{=} \frac{\lambda^k}{k!} e^{-\lambda}$$

# Algorithmic ride sharing



Probability of  $k$  requests from this area in the next 1 minute?

On average,  $\lambda = 5$  requests per minute

$$P(X = k) = \frac{\lambda^k}{k!} e^{-\lambda}$$

**Poisson  
distribution**



# Poisson Random Variable

Consider an experiment that lasts a fixed interval of time.

def A **Poisson** random variable  $X$  is the number of successes over the experiment duration, assuming **the time that each success occurs is independent** and the average # of requests over time is constant.

$$X \sim \text{Poi}(\lambda)$$

Support:  $\{0, 1, 2, \dots\}$

PMF

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

Expectation  $E[X] = \lambda$

Variance  $\text{Var}(X) = \lambda$

Examples:

- # earthquakes per year
- # server hits per second
- # of emails per day

Yes, expectation == variance  
for Poisson RV! More later!

# Earthquakes


$$X \sim \text{Poi}(\lambda) \quad p(k) = e^{-\lambda} \frac{\lambda^k}{k!}$$
$$E[X] = \lambda$$

There are an average of 2.79 major earthquakes in the world each year, and major earthquakes occur independently.

What is the probability of 3 major earthquakes happening next year?

1. Define RVs


2. Solve





# Other Discrete RVs

# Grid of random variables



# Geometric RV

Consider an experiment: independent trials of  $\text{Ber}(p)$  random variables.

def A **Geometric** random variable  $X$  is the # of trials until the first success.

$X \sim \text{Geo}(p)$

Support:  $\{1, 2, \dots\}$

PMF

$$P(X = k) = (1 - p)^{k-1} p$$

Expectation

$$E[X] = \frac{1}{p}$$

Variance

$$\text{Var}(X) = \frac{1-p}{p^2}$$

Examples:

- Flipping a coin ( $P(\text{heads}) = p$ ) until first heads appears
- Generate bits with  $P(\text{bit} = 1) = p$  until first 1 generated

# Negative Binomial RV

Consider an experiment: independent trials of  $\text{Ber}(p)$  random variables.

def A **Negative Binomial** random variable  $X$  is the # of trials until  $r$  successes.


$$\begin{array}{ll} X \sim \text{NegBin}(r, p) & \text{PMF} \\ & P(X = k) = \binom{k-1}{r-1} (1-p)^{k-r} p^r \\ & \text{Expectation} \\ & E[X] = \frac{r}{p} \\ & \text{Variance} \\ & \text{Var}(X) = \frac{r(1-p)}{p^2} \\ \text{Support: } \{r, r+1, \dots\} & \end{array}$$

Examples:

- Flipping a coin until  $r^{\text{th}}$  heads appears
- # of strings to hash into table until bucket 1 has  $r$  entries

$$\text{Geo}(p) = \text{NegBin}(1, p)$$

# Grid of random variables



# Catching Pokemon

Wild Pokemon are captured by throwing Pokeballs at them.

- Each ball has probability  $p = 0.1$  of capturing the Pokemon.
- Each ball is an independent trial.

What is the probability that you catch the Pokemon on the 5<sup>th</sup> try?


1. Define events/  
RVs & state goal

$X \sim$  some distribution

Want:  $P(X = 5)$

2. Solve

- A.  $X \sim \text{Bin}(5, 0.1)$
- B.  $X \sim \text{Poi}(0.5)$
- C.  $X \sim \text{NegBin}(5, 0.1)$
- D.  $X \sim \text{NegBin}(1, 0.1)$
- E.  $X \sim \text{Geo}(0.1)$






# Catching Pokemon

$$X \sim \text{Geo}(p) \quad p(k) = (1 - p)^{k-1}p$$

Wild Pokemon are captured by throwing Pokeballs at them.

- Each ball has probability  $p = 0.1$  of capturing the Pokemon.
- Each ball is an independent trial.

What is the probability that you catch the Pokemon on the 5<sup>th</sup> try?



1. Define events/  
RVs & state goal

$X \sim$  some distribution

Want:  $P(X = 5)$

2. Solve

- A.  $X \sim \text{Bin}(5, 0.1)$
- B.  $X \sim \text{Poi}(0.5)$
- C.  $X \sim \text{NegBin}(5, 0.1)$
- D.  $X \sim \text{NegBin}(1, 0.1)$
- E.  $X \sim \text{Geo}(0.1)$



# Exercises



The hardest part of is almost always deciding what you're modeling and what random variable to use.

# Kickboxing with RVs

How might you model the following?

1. # of snapchats you receive in a day
2. # of children born to the same parents until the first one with green eyes
3. If stock went up (1) or down (0) in a day
4. # of probability problems you try until you get 5 correct (if you are randomly correct)
5. # of years between now and 2050 with more than 6 Atlantic hurricanes

Choose from:

A. Ber( $p$ )	C. Poi( $\lambda$ )
B. Bin( $n, p$ )	D. Geo( $p$ )
	E. NegBin( $r, p$ )



# Kickboxing with RVs

How might you model the following?

1. # of snapchats you receive in a day
2. # of children born to the same parents until the first one with green eyes
3. If stock went up (1) or down (0) in a day
4. # of probability problems you try until you get 5 correct (if you are randomly correct)
5. # of years between now and 2050 with more than 6 Atlantic hurricanes

Note: These exercises are designed to build intuition; in a problem statement, you'll generally be given more detail.

Choose from: C. Poi( $\lambda$ )  
A. Ber( $p$ ) D. Geo( $p$ )  
B. Bin( $n, p$ ) E. NegBin( $r, p$ )

C. Poi( $\lambda$ )

D. Geo( $p$ ) or E. NegBin( $1, p$ )

A. Ber( $p$ ) or B. Bin( $1, p$ )

E. NegBin( $r = 5, p$ )

B. Bin( $n = 27, p$ ), where  
 $p = P(\geq 6 \text{ hurricanes in a year})$   
calculated from C. Poi( $\lambda$ )

# Poisson Random Variable

Review

$X \sim \text{Poi}(\lambda)$	PMF	$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$
	Expectation	$E[X] = \lambda$
Support: $\{0, 1, 2, \dots\}$	Variance	$\text{Var}(X) = \lambda$

In CS109, a Poisson RV  $X \sim \text{Poi}(\lambda)$  most often models

1. # of successes in a fixed interval of time, where successes are independent  
 $\lambda = E[X]$ , average success/interval

# 1. Web server load

$$X \sim \text{Poi}(\lambda) \quad p(k) = e^{-\lambda} \frac{\lambda^k}{k!}$$
$$E[X] = \lambda$$

Consider requests to a web server in 1 second.

- In the past, server load averages 2 hits/second, where requests arrive independently.
- Let  $X = \#$  requests the server receives in a second.

What is  $P(X < 5)$ ?

Define RVs

Solve

# Poisson Random Variable

$$X \sim \text{Poi}(\lambda)$$

PMF

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

Expectation  $E[X] = \lambda$

Support:  $\{0, 1, 2, \dots\}$

Variance  $\text{Var}(X) = \lambda$


In CS109, a Poisson RV  $X \sim \text{Poi}(\lambda)$  most often models

1. # of successes in a fixed time interval, where successes are independent  
 $\lambda = E[X]$ , average success/interval
2. **Approximation of  $Y \sim \text{Bin}(n, p)$**  where  $n$  is large and  $p$  is small.  
 $\lambda = E[Y] = np$

Approximation works well even when trials not entirely independent.

## 2. DNA

---



All the movies, images, emails and other digital data from more than 600 smartphones (10,000 GB) can be stored in the faint pink smear of DNA at the end of this test tube.

What is the probability that DNA storage stays uncorrupted?



## 2. DNA

What is the probability that DNA storage stays uncorrupted?


- In DNA (and real networks), we store large strings.
- Let string length be long, e.g.,  $n \approx 10^4$
- Probability of corruption of each base pair is very small, e.g.,  $p = 10^{-6}$
- Let  $X = \#$  of corruptions.

What is  $P(\text{DNA storage is uncorrupted}) = P(X = 0)$ ?

1. Approach 1:

$$X \sim \text{Bin}(n = 10^4, p = 10^{-6})$$

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$$


unwieldy!   $= \binom{10^4}{0} 10^{-6 \cdot 0} (1 - 10^{-6})^{10^4 - 0}$   
 $\approx 0.990049829$

2. Approach 2:

$$X \sim \text{Poi}(\lambda = 10^4 \cdot 10^{-6} = 0.01)$$

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!} = e^{-0.01} \frac{0.01^0}{0!}$$

$$= e^{-0.01}$$

$\approx 0.990049834$  a good approximation! 

# When is a Poisson approximation appropriate?

$$P(X = k) = \lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} = \dots$$

Def natural exponent

$$= \lim_{n \rightarrow \infty} \frac{n!}{n^k (n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

Expand

$$= \lim_{n \rightarrow \infty} \frac{n(n-1) \dots (n-k+1)}{n^k} \frac{(n-k)!}{(n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

Limit analysis

$$= \lim_{n \rightarrow \infty} \frac{n^k}{n^k} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{1}$$

Simplify

$$= \frac{\lambda^k}{k!} e^{-\lambda}$$

Under which conditions will  $X \sim \text{Bin}(n, p)$  behave like  $\text{Poi}(\lambda)$ , where  $\lambda = np$ ?

- A. Large  $n$ , large  $p$
- B. Small  $n$ , small  $p$
- C. Large  $n$ , small  $p$
- D. Small  $n$ , large  $p$
- E. Other



# Poisson approximation

$$X \sim \text{Poi}(\lambda)$$
$$E[X] = \lambda$$

$$Y \sim \text{Bin}(n, p)$$
$$E[Y] = np$$


Poisson approximates Binomial when  $n$  is large,  $p$  is small, and  $\lambda = np$  is "moderate".

Different interpretations of "moderate":

- $n > 20$  and  $p < 0.05$
- $n > 100$  and  $p < 0.1$

Poisson is Binomial in the limit:

- $\lambda = np$ , where  $n \rightarrow \infty, p \rightarrow 0$



# Poisson Random Variable

Consider an experiment that lasts a fixed interval of time.

def A **Poisson** random variable  $X$  is the number of occurrences over the experiment duration.

$$X \sim \text{Poi}(\lambda)$$

Support:  $\{0, 1, 2, \dots\}$

PMF

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

Expectation  $E[X] = \lambda$

Variance  $\text{Var}(X) = \lambda$

Examples:

- # earthquakes per year
- # server hits per second
- # of emails per day

Time to show intuition for why  
expectation == variance!

# Properties of $\text{Poi}(\lambda)$ with the Poisson paradigm

Recall the Binomial:

$$Y \sim \text{Bin}(n, p) \quad \begin{array}{ll} \text{Expectation} & E[Y] = np \\ \text{Variance} & \text{Var}(Y) = np(1 - p) \end{array}$$

Consider  $X \sim \text{Poi}(\lambda)$ , where  $\lambda = np$  ( $n \rightarrow \infty, p \rightarrow 0$ ):

$$X \sim \text{Poi}(\lambda) \quad \begin{array}{ll} \text{Expectation} & E[X] = \lambda \\ \text{Variance} & \text{Var}(X) = \lambda \end{array}$$

Proof:

$$E[X] = np = \lambda$$
$$\text{Var}(X) = np(1 - p) \rightarrow \lambda(1 - 0) = \lambda$$



# Poisson Approximation, approximately

---

Poisson can still provide a **good approximation of the Binomial**, even when assumptions are "mildly" violated.

You still can apply the Poisson approximation when:

- "Successes" in trials are almost, but not entirely independent e.g., # entries in each bucket in large hash table.
- Probability of "success" in each trial varies (slightly), like a **small relative change** in a very small  $p$  e.g., average # requests to web server/sec may fluctuate slightly due to load on network or time of day



We won't explore this too much, but we want you to know about it anyway.

# Can these Binomial RVs be approximated?




Poisson approximates Binomial when  $n$  is large,  $p$  is small, and  $\lambda = np$  is "moderate".

Different interpretations of "moderate":

- $n > 20$  and  $p < 0.05$
- $n > 100$  and  $p < 0.1$

Poisson is Binomial in the limit:

- $\lambda = np$ , where  $n \rightarrow \infty, p \rightarrow 0$



# Can these Binomial RVs be approximated?

Poisson approximates Binomial when  $n$  is large,  $p$  is small, and  $\lambda = np$  is "moderate".

Different interpretations of "moderate":

- $n > 20$  and  $p < 0.05$
- $n > 100$  and  $p < 0.1$

Poisson is Binomial in the limit:

- $\lambda = np$ , where  $n \rightarrow \infty, p \rightarrow 0$

